



FIG. 1




FIG. 2A

Flag-Smad2	-	WT	EPSM	3S-A
TGF-β ₁	-	- +	- +	- +



FIG. 2B BEST AVAILABLE COPY





FIG. 2C



BEST AVAILABLE COPY

FIG. 3





FIG. 4


BEST AVAILABLE COPY



(MEAN \pm S.D. n=7)

FIG. 5A


DETECTION OF LOCALIZATION IN CELLS



BEST AVAILABLE COPY

FIG. 5B


DETECTION OF INFLUENCE BY INHIBITOR




BEST AVAILABLE COPY

FIG. 6

IMMUNOHISTOLOGICAL EXAMINATION OF PHOSPHORYLATION AT
Smad3 LINKER REGION IN LIVER OF HEPATIC FIBROSIS MODEL RAT



pSmad3L









αSMA

BEST AVAILABLE COPY

FIG. 7

KINASE ASSAY
 IP: α pp38

TGF- β_1		-	+
GST-Smad2	IB: α pSmad2L		- MAPK - dependent pSmad2
	α pSmad2C		- T β RI - dependent pSmad2
	α Smad2/3		- GST-Smad2
GST-Smad3	α pSmad3L		- MAPK - dependent pSmad3
	α pSmad3C		- T β RI - dependent pSmad3
	α Smad2/3		- GST-Smad3

BEST AVAILABLE COPY

FIG. 8

α pSmad3L



α Ki-67

